

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
19 July 2001 (19.07.2001)

PCT

(10) International Publication Number
WO 01/51454 A1

(51) International Patent Classification⁷: C07C 229/00,
A61K 38/28, 51/00

Doris, C. [US/US]; 33 Lakeview Road, Carmel, NY 10512 (US).

(21) International Application Number: PCT/US01/01274

(74) Agents: LESSLER, Jay, P. et al.; Darby & Darby P.C.,
805 Third Avenue, New York, NY 10022-7513 (US).

(22) International Filing Date: 12 January 2001 (12.01.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/175,947 13 January 2000 (13.01.2000) US
60/194,421 4 April 2000 (04.04.2000) US
60/202,210 5 May 2000 (05.05.2000) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): EMI-SPHERE TECHNOLOGIES, INC. [US/US]; 765 Old Saw Mill River Road, Tarrytown, NY 10591 (US).

Published:

— with international search report

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): BOYD, Maria, Aurora, P. [PH/US]; 43 South Highland Avenue, Ossining, NY 10562 (US). LEONE-BAY, Andrea [US/US]; 20 Woodland Way, Ridgefield, CT 06877 (US). O'TOOLE,

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/51454 A1

(54) Title: COMPOUNDS AND COMPOSITIONS FOR DELIVERING ACTIVE AGENTS

(57) Abstract: Compounds and compositions for the delivery of active agents are provided. Methods of administration and preparation are provided as well.

5 **Compounds And Compositions For Delivering Active Agents**

FIELD OF THE INVENTION

10 The present invention relates to compounds for delivering
active agents, such as biologically or chemically active agents,
to a target. These compounds are well suited for forming non-
covalent mixtures with active agents for oral, intracolonic,
pulmonary, and other routes of administration to animals.
Methods for the preparation and administration of such
15 compositions are also disclosed.

BACKGROUND OF THE INVENTION

20 Conventional means for delivering active agents are often
severely limited by biological, chemical, and physical barriers.

Typically, these barriers are imposed by the environment
through which delivery occurs, the environment of the target for
delivery, and/or the target itself. Biologically and chemically
active agents are particularly vulnerable to such barriers.

25 In the delivery to animals of biologically active and
chemically active pharmacological and therapeutic agents,
barriers are imposed by the body. Examples of physical barriers
are the skin, lipid bi-layers and various organ membranes that
are relatively impermeable to certain active agents but must be
30 traversed before reaching a target, such as the circulatory
system. Chemical barriers include, but are not limited to, pH
variations in the gastrointestinal (GI) tract and degrading
enzymes.

These barriers are of particular significance in the design of oral delivery systems. Oral delivery of many biologically or chemically active agents would be the route of choice for administration to animals if not for biological, chemical, and physical barriers. Among the numerous agents which are not typically amenable to oral administration are biologically or chemically active peptides, such as calcitonin and insulin; polysaccharides, and in particular mucopolysaccharides including, but not limited to, heparin; heparinoids; antibiotics; and other organic substances. These agents may be rapidly rendered ineffective or destroyed in the gastrointestinal tract by acid hydrolysis, enzymes, and the like. In addition, the size and structure of macromolecular drugs may prohibit absorption.

Earlier methods for orally administering vulnerable pharmacological agents have relied on the co-administration of adjuvants (e.g., resorcinols and non-ionic surfactants such as polyoxyethylene oleyl ether and n-hexadecylpolyethylene ether) to increase artificially the permeability of the intestinal walls, as well as the co-administration of enzymatic inhibitors (e.g., pancreatic trypsin inhibitors, diisopropylfluorophosphate (DFF) and trasylol) to inhibit enzymatic degradation. Liposomes have also been described as drug delivery systems for insulin and heparin. However, broad spectrum use of such drug delivery systems is precluded because: (1) the systems require toxic amounts of adjuvants or inhibitors; (2) suitable low molecular weight cargos, i.e. active agents, are not available; (3) the systems exhibit poor stability and inadequate shelf life; (4) the systems are difficult to manufacture; (5) the systems fail to protect the active agent (cargo); (6) the systems adversely alter the active agent; or (7) the systems fail to allow or

promote absorption of the active agent.


More recently, proteinoid microspheres have been used to deliver pharmaceuticals. See, for example, US Patent Nos. 5,401,516; 5,443,841; and Re. 35,862. In addition, certain
5 modified amino acids have been used to deliver pharmaceuticals.

See, for example, US Patent Nos. 5,629,020; 5,643,957; 5,766,633; 5,776,888; and 5,866,536.

However, there is still a need for simple, inexpensive delivery systems which are easily prepared and which can deliver
10 a broad range of active agents by various routes.

SUMMARY OF THE INVENTION

The present invention provides compounds and compositions which facilitate the delivery of active agents. Delivery agent
15 compounds of the present invention have the following formula:



Formula I

wherein:

20 R^1 , R^2 , R^3 , R^4 , and R^5 are independently selected from hydrogen, halogen, $-\text{OH}$, $-\text{OCH}_3$, $\text{C}_1\text{-C}_4$ alkyl, $-\text{NH}_2$, $-\text{NH}(\text{CH}_3)$, $-\text{N}(\text{CH}_3)_2$, or $-\text{NO}_2$;

m is an integer ranging from 0 to 4;

25 R^6 is a phenyl substituted with $-\text{O}-\text{R}^7\text{-COOH}$ at the ortho, meta, or para position;

R^6 is optionally substituted with one or more members selected from halogen, $-\text{OH}$, $-\text{OCH}_3$, $\text{C}_1\text{-C}_4$ alkyl, $-\text{NH}_2$, $\text{NH}(\text{CH}_3)$, -

$N(CH_3)_2$, and $-NO_2$; and

R^7 is C_1 - C_{12} alkyl,

or a salt thereof.

In one preferred embodiment, R^1 is $-OH$. According to
 5 another preferred embodiment, R^1 is $-OCH_3$.

In another preferred embodiment, m is 0. According to yet
 another preferred embodiment, m is 3.

In another preferred embodiment, the phenyl group in R^6 is
 substituted with Cl, preferably para to the $-O-R^7-COOH$.

10 In another preferred embodiment, R^7 is C_3 - C_9 alkyl.


According to yet another preferred embodiment, R^7 is C_3 - C_7
 alkyl.

In another preferred embodiment, R^1 is $-OH$, R^6 is a phenyl,
 $-O-R^7-COOH$ in the ortho position. More preferably, R^6 is a
 15 phenyl, with a Cl substituent para to the $-O-R^7-COOH$. . More
 preferably, R^7 is a C_3 - C_7 alkyl. More preferably, $m = 0$.

In another preferred embodiment, R^1 is $-OCH_3$, m is from 1
 to 3, R^6 is a phenyl substituted with $-O-R^7-COOH$ at the para
 position. More preferably, R^7 is a C_3 - C_7 alkyl.

20 Preferred compounds include, but are not limited to, the
 monosodium and disodium salts of the compounds of formula I
 above.

Special mention is made of compounds having the formula:



25

Formula II

wherein

n is an integer from 1 to 12;

X is one or more of the following independently selected
5 substituents: halogen, -OH, -OCH₃, C₁-C₄ alkyl, -NH₂, -N(CH₃)₂,
or -NO₂;


Y is one or more of the following independently selected
substituents: halogen, OH, OCH₃, C₁-C₄ alkyl, NH₂, N(CH₃)₂, or NO₂,
or salts thereof.

10 Preferably, n is an integer from 3 to 9 and more preferably
from 3 to 7.

According to one preferred embodiment, X is a Cl at the 4-
position of the phenyl group.


According to another preferred embodiment, Y is -OH at the
15 2-position of the phenyl group.

More preferred compounds comprise one or more of the
following:




20

Compound 1




Compound 2




Compound 3

5








Compound 4



Compound 5

10

Compound 6Compound 7Compound 8Compound 9

Compound 10

5 or salts thereof.

The invention also provides a composition comprising at least one of the delivery agent compounds of the formulae I and II, 1-10, above, or salts thereof, or mixtures thereof, and at least one active agent. These compositions deliver active
10 agents to biological systems in increased or improved bioavailability of the active agent compared to administration of the active agent without the delivery agent compound.

Also provided are dosage unit forms comprising the compositions. The dosage unit form may be in the form of a
15 liquid or a solid, such as a tablet, capsule or particle, including a powder or sachet.

Another embodiment is a method for administering an active agent to an animal in need of the active agent by administering to the animal a composition comprising at least one of the
20 delivery agent compounds above and the active agent. Preferred routes of administration include the oral, intracolonic and pulmonary routes.

Yet another embodiment is a method of treating a disease or for achieving a desired physiological effect in an animal by
25 administering the composition of the present invention.

Yet another embodiment is a method of preparing a composition of the present invention by mixing at least one delivery agent compound of the formula above and at least one

active agent.

DETAILED DESCRIPTION OF THE INVENTION

5

Delivery Agent Compounds

The delivery agent compounds may be in the form of the carboxylic acid or salts thereof. Suitable salts include, but are not limited to, organic and inorganic salts, for example
10 alkali-metal salts, such as sodium, potassium and lithium; alkaline-earth metal salts, such as magnesium, calcium or barium; ammonium salts; basic amino acids, such as lysine or arginine; and organic amines, such as dimethylamine or pyridine.

Preferably, the salts are sodium salts. The salts may be mono-
15 or multi-valent salts, such as monosodium salts and di-sodium salts. The salts may also be solvates, including ethanol solvates, and hydrates. The term "solvate" as used herein includes, but is not limited to, a molecular or ionic complex of molecules or ions of a solvent, such as ethanol, with ions or
20 molecules of the compounds of the present invention.

Salts of the delivery agent compounds of the present invention may be prepared by methods known in the art. For example, sodium salts may be prepared by dissolving the delivery agent compound in ethanol and adding aqueous sodium hydroxide.

25 In general, the compounds of the present invention may be prepared using the following method.

The appropriately substituted amino-phenol (such as 2-amino-phenol) and the appropriate alkylating agent are mixed in potassium hydroxide (KOH) and dimethyl sulfoxide (DMSO) for 2
30 hours at room temperature, then poured into distilled water and extracted with ethyl acetate (EtOAc). The combined organic

layers are washed with water, dried with magnesium sulfate (MgSO_4), and filtered. The solvent is evaporated *in vacuo* to give a crude product, which may be purified by methods known in the art, such as medium pressure liquid chromatography (MPLC) utilizing 10% EtOAc in hexanes as the eluting solvent, to yield an intermediate ester.

The intermediate ester is dissolved in tetrahydrofuran (THF), flushed with nitrogen and cooled to about 0°C. Triethylamine and then the appropriately substituted benzoyl chloride dissolved in THF (100 mL) are added slowly to the solution containing the intermediate ester. The mixture is allowed to warm up to room temperature and is stirred overnight, monitoring the progress of the reaction by thin-layer chromatography. Work-up of the reaction is done by addition of 3% HCl solution. The phases are separated, and the organic layer is washed with 3% HCl, water and then a saturated solution of NaCl. The organic layer is dried with MgSO_4 , filtered, and the solvent evaporated *in vacuo*.

A 10% aqueous solution of sodium hydroxide (NaOH) is added to this product and the mixture stirred for 2 hours at about 90°C, then cooled and diluted with water. The solution is acidified to about a pH of 6 or lower. The product separates from solution and is collected by filtration, washed generously with water, and dried overnight under vacuum at 40°C to yield the product as a crystalline powder.

The term "halogen" as used herein, unless otherwise indicated, fluoro, chloro, bromo and iodo. The term "alkyl" as used herein, unless otherwise indicated, includes saturated monovalent hydrocarbon radicals and may be straight, branched or cyclic. Such alkyl groups may include one or two double or triple bonds. It is understood that for cyclic alkyls at least

three carbon atoms are required to form the ring.

The solvate may be prepared by dissolving the compound of the present invention in a solvent, such as ethanol, and then reacting the mixture with a molar excess of a salt, such as a sodium containing salt (e.g. sodium hydroxide). For example, to
5 prepare an ethanol solvate of a disodium salt of a compound of the present invention, at least about two molar equivalents of a monosodium containing salt are added to a mixture of the compound and ethanol. The solvate may be recovered by methods
10 known in the art.

Active Agents

Active agents suitable for use in the present invention include biologically active agents and chemically active agents,
15 including, but not limited to, pesticides, pharmacological agents, and therapeutic agents.

For example, biologically or chemically active agents suitable for use in the present invention include, but are not limited to, proteins; polypeptides; peptides; hormones;
20 polysaccharides, and particularly mixtures of muco-polysaccharides; carbohydrates; lipids; small polar organic molecules (i.e. polar organic molecules having a molecular weight of 500 daltons or less); other organic compounds; and particularly compounds which by themselves do not pass (or which
25 pass only a fraction of the administered dose) through the gastro-intestinal mucosa and/or are susceptible to chemical cleavage by acids and enzymes in the gastro-intestinal tract; or any combination thereof.

Further examples include, but are not limited to, the
30 following, including synthetic, natural or recombinant sources thereof: growth hormones, including human growth hormones

(hGH), recombinant human growth hormones (rhGH), bovine growth hormones, and porcine growth hormones; growth hormone releasing hormones; growth hormone releasing factor, interferons, including α , β and γ ; interleukin-1; interleukin-2; insulin, including porcine, bovine, human, and human recombinant, optionally having counter ions including zinc, sodium, calcium and ammonium; insulin-like growth factor, including IGF-1; heparin, including unfractionated heparin, heparinoids, dermatans, chondroitins, low molecular weight heparin, very low molecular weight heparin and ultra low molecular weight heparin; calcitonin, including salmon, eel, porcine and human; erythropoietin; atrial natriuretic factor; antigens; monoclonal antibodies; somatostatin; protease inhibitors; adrenocorticotropin, gonadotropin releasing hormone; oxytocin; leutinizing-hormone-releasing-hormone; follicle stimulating hormone; glucocerebrosidase; thrombopoietin; filgrastim; prostaglandins; cyclosporin; vasopressin; cromolyn sodium (e.g. sodium or disodium chromoglycate); vancomycin; desferrioxamine (DFO); bisphosphonates, including alendronate, tiludronate, etidronate, clodronate, pamidronate, olpadronate, and incadronate; parathyroid hormone, including its fragments; antimicrobials, including antibiotics, anti-bacterials (non-limiting examples of antibiotics include gram-positive acting, bacteriocidal, lipopeptidal and cyclic peptidal antibiotics, such as daptomycin and analogs thereof) and anti-fungal agents; vitamins; analogs, fragments, mimetics or polyethylene glycol (PEG)-modified derivatives of these compounds; or any combination thereof.

Delivery systems

The composition of the present invention comprises one or more delivery agent compounds of the present invention, and one or more active agents. In one embodiment, one or more of the
5 delivery agent compounds, or salts of these compounds, or poly amino acids or peptides of which these compounds or salts form one or more of the units thereof, may be used as a delivery agent by mixing with the active agent prior to administration to form an administration composition.

10 The administration compositions may be in the form of a liquid. The solution medium may be water (for example, for salmon calcitonin, parathyroid hormone, and erythropoietin), 25% aqueous propylene glycol (for example, for heparin) and phosphate buffer (for example, for rhGH). Other dosing vehicles
15 include polyethylene glycol. Dosing solutions may be prepared by mixing a solution of the delivery agent compound with a solution of the active agent, just prior to administration. Alternatively, a solution of the delivery agent compound (or active agent) may be mixed with the solid form of the active
20 agent (or delivery agent compound). The delivery agent compound and the active agent may also be mixed as dry powders. The delivery agent compound and the active agent can also be admixed during the manufacturing process.

The dosing solutions may optionally contain additives such
25 as phosphate buffer salts, citric acid, glycols, or other dispersing agents. Stabilizing additives may be incorporated into the solution, preferably at a concentration ranging between about 0.1 and 20% (w/v).

The administration compositions may alternately be in the
30 form of a solid, such as a tablet, capsule or particle, such as a powder or sachet. Solid dosage forms may be prepared by

mixing the solid form of the compound with the solid form of the active agent. Alternately, a solid may be obtained from a solution of compound and active agent by methods known in the art, such as freeze-drying (lyophilization), precipitation, crystallization and solid dispersion.

The administration compositions of the present invention may also include one or more enzyme inhibitors. Such enzyme inhibitors include, but are not limited to, compounds such as actinonin or epiactinonin and derivatives thereof. Other enzyme inhibitors include, but are not limited to, aprotinin (Trasylol) and Bowman-Birk inhibitor.

The amount of active agent used in an administration composition of the present invention is an amount effective to accomplish the purpose of the particular active agent for the target indication. The amount of active agent in the compositions typically is a pharmacologically, biologically, therapeutically, or chemically effective amount. However, the amount can be less than that amount when the composition is used in a dosage unit form because the dosage unit form may contain a plurality of delivery agent compound/active agent compositions or may contain a divided pharmacologically, biologically, therapeutically, or chemically effective amount. The total effective amount can then be administered in cumulative units containing, in total, an effective amount of the active agent.

The total amount of active agent to be used can be determined by methods known to those skilled in the art. However, because the compositions of the invention may deliver active agents more efficiently than compositions containing the active agent alone, lower amounts of biologically or chemically active agents than those used in prior dosage unit forms or delivery systems can be administered to the subject, while still

achieving the same blood levels and/or therapeutic effects.

The presently disclosed delivery agent compounds facilitate the delivery of biologically and chemically active agents, particularly in oral, intranasal, sublingual, intraduodenal, subcutaneous, buccal, intracolonic, rectal, vaginal, mucosal, pulmonary, transdermal, intradermal, parenteral, intravenous, intramuscular and ocular systems, as well as traversing the blood-brain barrier.

Dosage unit forms can also include any one or combination of excipients, diluents, disintegrants, lubricants, plasticizers, colorants, flavorants, taste-masking agents, sugars, sweeteners, salts, and dosing vehicles, including, but not limited to, water, 1,2-propane diol, ethanol, olive oil, or any combination thereof. Non-limiting examples of such ingredients are described in Remingtons Pharmaceutical Sciences, 18th Ed., Mack Publishing Co. (1990), hereby incorporated by reference.

The compounds and compositions of the subject invention are useful for administering biologically or chemically active agents to any animals, including but not limited to birds, such as chickens; mammals, such as rodents, cows, pigs, dogs, cats, primates, and particularly humans; and insects.

The system is particularly advantageous for delivering chemically or biologically active agents that would otherwise be destroyed or rendered less effective by conditions encountered before the active agent reaches its target zone (i.e. the area in which the active agent of the delivery composition is to be released) and within the body of the animal to which they are administered. Particularly, the compounds and compositions of the present invention are useful in orally administering active agents, especially those that are not ordinarily orally deliverable, or those for which improved delivery is desired.

The compositions comprising the compounds and active agents have utility in the delivery of active agents to biological systems and in an increased or improved bioavailability of the active agent compared to administration of the active agent without the delivery agent. Delivery can be improved by delivering more active agent over a period of time, or in delivering active agent in a particular time period (such as to effect quicker or delayed delivery) or over a period of time (such as sustained delivery).

Another embodiment of the present invention is a method for the treatment or prevention of a disease or for achieving a desired physiological effect, such as those listed in the table below, in an animal by administering the composition of the present invention. Specific indications for active agents can be found in the Physicians' Desk Reference (54th Ed., 2000, Medical Economics Company, Inc., Montvale, NJ), which is herein incorporated by reference. The active agents in the table below include their analogs, fragments, mimetics, and polyethylene glycol-modified derivatives.

Active Agent	Disease and Physiological Effect
Growth hormones	Growth disorders
Interferons, including α , β and γ	Viral infection, including chronic cancer and multiple sclerosis
Interleukin-1; interleukin-2	Viral infection; cancer
Insulin; Insulin-like growth factor (IGF-1)	Diabetes
Heparin	Thrombosis; prevention of blood coagulation
Calcitonin	Osteoporosis; diseases of the bone
Erythropoietin	Anemia
Atrial natriuretic factor	Vasodilation
Antigens	Infection
Monoclonal antibodies	To prevent graft rejection; cancer

Active Agent	Disease and Physiological Effect
Somatostatin	Bleeding ulcer; erosive gastritis
Protease inhibitors	AIDS
Adrenocorticotropin	High cholesterol (to lower cholesterol)
Gonadotropin releasing hormone	Ovulatory disfunction (to stimulate ovulation)
Oxytocin	Labor disfunction (to stimulate contractions)
Leutinizing-hormone-releasing-hormone; follicle stimulating hormone	Regulate reproductive function
Glucocerebrosidase	Gaucher disease (to metabolize lipoprotein)
Thrombopoietin	Thrombocytopenia
Filgrastim	Reduce infection in chemotherapy patients
Prostaglandins	Hypertension
Cyclosporin	Transplant rejection
Vasopressin	Bed-wetting; antidiuretic
Cromolyn sodium; Vancomycin	Asthma; allergies
Desferrioxamine (DFO)	Iron overload
Parathyroid hormone (PTH), including its fragments	Osteoporosis; Diseases of the bone
Antimicrobials	Infection including gram-positive bacterial infection
Vitamins	Vitamin deficiencies
Bisphosphonates	Osteoporosis ; Paget's disease; Inhibits osteoclasts

For example, one embodiment of the present invention is a method for treating a patient suffering from or susceptible to diabetes by administering insulin and at least one of the delivery agent compounds of the present invention.

Following administration, the active agent present in the composition or dosage unit form is taken up into the circulation. The bioavailability of the agent is readily assessed by measuring a known pharmacological activity in blood, e.g. an increase in blood clotting time caused by heparin, or a

decrease in circulating calcium levels caused by calcitonin. Alternately, the circulating levels of the active agent itself can be measured directly.

5 DESCRIPTION OF THE PREFERRED EMBODIMENTS

The following examples illustrate the invention without limitation. All parts are given by weight unless otherwise indicated.

Proton nuclear magnetic resonance (^1H NMR) analyses for the
10 compounds listed below were conducted on a 300 MHz Bruker spectrometer using dimethyl sulfoxide (DMSO-d_6) as the solvent unless otherwise indicated.

Example 1 - Compound Preparation

15 Preparation of compound 1

A 500-mL round-bottomed flask was charged with KOH (16.1 g, 270 mmol) and DMSO (160 mL). 2-Amino-4-chlorophenol (13.6 g, 94.7 mmol) was added to the mixture, followed by ethyl 8-bromooctanoate (24.0 g, 95.5 mmol), and the resulting mixture
20 was stirred for 2 hours at room temperature. The mixture was poured into 200 mL of distilled water. The solution was extracted with EtOAc (2 X 200 mL). The organic layers were combined, washed once with water, dried with MgSO_4 , and filtered.

The solvent was evaporated *in vacuo* to give a crude product,
25 which was a thick, brown liquid. The product was purified by medium pressure liquid chromatography (MPLC) utilizing 10% EtOAc in hexanes as the eluting solvent, to yield an intermediate ester as a light brown liquid (17.1 g, 54.5 mmol, 38% yield).

The intermediate ester (3.37 g, 17 mmol) was dissolved in
30 THF (150 mL) and charged to a 500-mL round-bottomed flask equipped with an addition funnel. The system was flushed with

nitrogen and cooled to 0°C with an ice bath. Triethylamine (3.0 mL, 21.2 mmol) was added slowly via a syringe. O-Acetyl salicyloyl chloride (3.37 g, 17 mmol) was dissolved in THF (100 mL) and charged to the addition funnel and added slowly over 30 minutes to the ester mixture. The solution slowly turned cloudy. The mixture was allowed to warm up to room temperature and was stirred overnight. The progress of the reaction was monitored by thin-layer chromatography. Work-up of the reaction was done by addition of 3% (v/v) HCl solution (100 mL). The phases were separated, and the organic layer was washed once more with 3% (v/v) HCl (100 mL), followed by H₂O (100 mL) and a saturated solution of NaCl (100 mL). The organic layer was dried with MgSO₄, filtered, and the solvent was evaporated *in vacuo*.

A 10% (w/w) aqueous solution of NaOH (68 mL) was added to the crude product isolated above and the mixture was stirred for 2 hours at 90°C using an oil bath. The solution was cooled and diluted with water (100 mL), and acidified to a pH of about 6. A white precipitate was formed, collected by filtration, washed generously with water, and dried overnight under vacuum at 40°C to yield **Compound 1** as a fine, white crystalline powder (4.60 g, 11.3 mmol, 66% yield); mp 152-153°C; Analysis calculated for C₂₁H₂₄ClNO₅: %C, 62.14 (calculated), 62.20 (found); %H, 5.96 (calculated), 5.81 (found); %N, 3.45 (calculated), 3.26 (found). ¹H NMR Analysis (DMSO-d₆): δ 1.27-1.53, 1.77-1.91, 2.15-2.20, (4.05-4.09, 6.97-7.10, 7.41-7.47, 8.01-8.05, 8.56-8.57, 11.0, 11.6, 12.0. Compounds 2, 3, 4, and 5 were prepared by this method using the appropriate starting materials. Compound 6 can also be prepared by this method using the appropriate starting materials.

Compound 2 Melting point: 166-169° C. Combustion analysis:

%C: 59.43 (calculated), 59.37 (found); %H: 4.99 (calculated), 4.96 (found); %N: 3.85 (calculated), 3.59 (found); %Cl: 9.75 (calculated), 9.74 (found). ¹H NMR Analysis (d6-DMSO): δ 11.80-12.0, 11.00, 8.58, 8.02-8.05, 7.41-7.47, 7.06-7.10, 6.98-7.03 ,
5 4.06-4.11, 2.31-2.36 , 1.68-1.86.)

Compounds 3: Melting point: 231-234° C. Combustion analysis: %C: 58.38 (calculated), 58.13 (found); %H: 4.61 (calculated), 4.67 (found); %N: 4.00 (calculated), 3.91 (found); %Cl: 10.14 (calculated), 10.10 (found). ¹H NMR Analysis: (d6-DMSO): δ
10 11.80-12.0, 11.00 , 8.02-8.06, 7.41-7.47, 7.06-7.10, 6.98-7.03, 4.08-4.13 , 2.44-2.52, 2.00-2.10.

Compound 4: Melting point: 122-124 C. Combustion analysis: %C: 67.91 (calculated), 67.77 (found); %H: 6.78 (calculated), 6.83 (found); %N: 3.77 (calculated), 3.78 (found). ¹H NMR
15 Analysis: (d6-DMSO): δ 11.80-12.0 (br. s, 1 H), 11.00 (s, 1 H), 8.41-8.44 (m, 1 H), 8.01-8.04 (dd, 1 H), 7.38-7.44 (m, 1 H), 7.04-7.08 (m, 3 H), 6.91-7.00 (m, 2 H), 4.03-4.08 (t, 2 H), 2.15-2.19 (t, 2 H), 1.78-1.83 (m, 2 H), 1.26-1.50 (m, 8 H).

Compound 5: Melting point: 147-149 C. Combustion analysis: 20 %C: 54.22 (calculated), 54.15 (found); %H: 5.79 (calculated), 5.74 (found); %N: 5.75 (calculated), 5.66 (found). ¹H NMR Analysis: (d6-DMSO): δ 11.80-12.0 (br. s, 1 H), 11.00 (s, 1 H), 8.44-8.47 (m, 1 H), 8.02-8.05 (dd, 1 H), 7.39-7.45 (m, 1 H), 7.01-7.09 (m, 3 H), 6.91-6.99 (m, 2 H), 4.05-4.10 (t, 2 H),
25 2.31-2.36 (t, 2 H), 1.68-1.87 (m, 4 H).

Preparation of Compound 8

Step A: 4-aminophenol (10.00g, 0.0917), dioxane (175 ml), and water (90 ml) were added to a 1 liter, round, bottomed flask
30 fitted with a magnetic stir bar, addition funnel and argon purge. The reaction mixture was cooled in an ice/water bath and

dropwise addition of a 1 N NaOH solution (91.7 ml) was made to the reaction mixture over a period of 10 minutes. Di-*t*-butyl dicarbonate (19.00g, 0.0917 mol) was dissolved in dioxane (30 ml) and this solution was placed in the addition funnel.

5 Dropwise addition of this solution was made over a 45 minute period. The ice bath was removed and the reaction mixture was allowed to come to ambient temperature overnight. After stirring overnight, dioxane was removed *in vacuo* and ethyl acetate was added to the reaction mixture. 0.5M H₂SO₄ solution
10 was added until the pH of the aqueous layer was 2. The ethyl acetate layer was separated off and the aqueous layer was extracted with two additional portions of ethyl acetate. The combined ethyl acetate layers were decolorized with activated carbon, were dried with sodium sulfate and concentrated *in vacuo*
15 to yield a brown solid which was washed with warm hexanes. The resulting 4-N-(*tert*-butyloxycarbonyl)-1-hydroxy benzene was isolated in a yield of 14.82 g.

A portion of the 4-N-(*tert*-butyloxycarbonyl)-1-hydroxy benzene (7.0 g, 0.0335 mol) and acetone (60 ml) were placed in a
20 250 ml round bottomed flask fitted with argon purge, magnetic stir bar and cold water condenser. 3-bromobenzylpropyl ether (6.50 ml, 0.0368 mol) and potassium carbonate (4.62 g, 0.0335 mol) were added and heating of the reaction mixture was started.

After refluxing for 4 days, heating of the reaction mixture was
25 stopped and deionized water (100 ml) was added. This mixture was extracted with 2 portions of diethyl ether (100 ml portions). The combined ether layer was then extracted with 2 portions of 2N NaOH solution (100 ml portions). The ether layer was dried with sodium sulfate and concentrated *in vacuo* leaving
30 a tan solid. The tan solid was recrystallized from hexanes to

yield 3-(4-N-(tert-butyloxycarbonyl)phenoxy)-1-benzyloxy propane (8.2 g).

3-(4-N-(tert-butyloxycarbonyl)phenoxy)-1-benzyloxy propane (12.75 g, 0.0357 mol), prepared as written above, was placed in a 500 ml round bottomed flask fitted with an argon purge and a magnetic stir bar. The reaction mixture was sparged with argon for about 30 minutes before 10% palladium on activated carbon (0.75 g) was added to the reaction mixture. The reaction vessel was evacuated and a balloon containing hydrogen was placed atop the reaction vessel. After stirring two days an additional portion of 10% palladium on activated carbon (0.25 g) was added.

The flask was again evacuated and a balloon containing hydrogen again was placed atop the flask. This mixture stirred for 4 additional days under hydrogen (refilling the balloon as necessary). The reaction mixture was then filtered and the filtrate was concentrated *in vacuo* yielding 3-(4-N-(tert-butyloxycarbonyl)phenoxy)propane-1-ol as a white solid in a yield of 9.47 g.

3-(4-N-(tert-butyloxycarbonyl)phenoxy)propane-1-ol (9.25 g, 0.0346 mol), which was prepared as described above, was added to a 250 ml, round, bottomed flask fitted with an argon purge and a magnetic stir bar. Pyridinium dichromate (45.59 g, 0.1213 mol) was added and the reaction was allowed to stir at ambient temperature overnight. The reaction mixture was then poured into 500 ml of water and was extracted with three, 200 ml portions of ethyl acetate. The combined ethyl acetate portions were washed with 100 ml of 0.5M H₂SO₄, and then were extracted with 100 ml of 2N, NaOH solution. The NaOH layer was then acidified with 2N HCl solution, which caused a gray solid to precipitate. 3-(4-N-(tert-butyloxycarbonyl)phenoxy)propionic acid was isolated in a yield of 3.26 g.

3-(4-N-(tert-butyloxycarbonyl)phenoxy)propionic acid prepared above (3.25 g, 0.01157 mol) and dioxane (30 ml) were placed in a 100 ml round, bottomed flask. The reaction mixture was filtered and the filtrate was placed in another 100 ml round
5 bottomed flask fitted with an argon purge, and magnetic stir bar. 4M HCl in dioxane (20 ml) was added and the mixture was allowed to stir at ambient temperature. After stirring overnight, the reaction mixture was filtered and a white solid was isolated. The filtrate was set-aside for three days and
10 more solid was noted to have precipitated. This solid was isolated by filtration and was combined with the earlier isolated solid. 4M HCl in dioxane (10 ml) was added to the solids. After stirring overnight at ambient temperature, additional solid was noted to have precipitated. This was also
15 isolated by filtration and combined with the earlier isolated solids. This combined solid was dried in vacuo and 3-(4-aminophenoxy)propionic acid was isolated in a yield of 2.08 g.

3-(4-aminophenoxy)propionic acid prepared above (1.95 g, 0.0108 mol) was placed in a 250 ml round bottomed flask fitted
20 with a condenser, argon purge, and magnetic stir bar. Dichloromethane was added and a tan slurry was formed. Trimethylsilyl chloride (2.74 ml, 0.0215 mol) was added to the reaction mixture and heating to reflux was started. After 1.5 hours, heating was discontinued and the reaction mixture was
25 cooled in an ice/water bath. Triethyl amine (2.25 ml, 0.0162 mol) was added. Acetylsalicyloyl chloride (2.03 g, 0.0102 mol) was dissolved in dichloromethane (10 ml). This solution was placed in an addition funnel atop the flask, which had replaced the condenser atop the flask. Dropwise addition of the
30 acetylsalicyloyl chloride solution was then made over a period of 15 minutes. This mixture was allowed to stir an additional

30 minutes before the ice bath was removed and the reaction was allowed to come to room temperature overnight. The methylene chloride was then removed in vacuo. 2N NaOH (70 ml) was then added and this mixture was allowed to stir for approximately 30 minutes before the reaction mixture was acidified with 2M H₂SO₄.

A tan solid precipitated which was isolated by filtration, taken up in ethyl acetate and washed with 2, 50 ml portions of 2N HCl solution. The ethyl acetate layer was dried with sodium sulfate and was concentrated in vacuo. The resulting tan solid was recrystallized from ethanol/water solution, washed with water resulting in a tan solid.

Step B: This solid was taken up in ethyl alcohol in a 250 ml Erlenmeyer flask. The mixture was heated until the solid was dissolved. The mixture was then hot filtered and 10N NaOH solution (0.64 ml, 0.0064 mol) was added to the filtrate to form a filtrate. After stirring the mixture for several minutes, approximately half of the ethanol was removed in vacuo. Heptane (75 ml) was added to the resulting slurry. The mixture was filtered, dried further in vacuo and a solid was isolated. The solid was taken up in a NaOH/H₂O mixture. This solution was acidified with 2M H₂SO₄. The white solid which precipitated was isolated by filtration, recrystallized from a 1:1, ethanol:water mixture and was dried under vacuum.

Step C: The product, 3-(4-salicyloylamino)phenoxy propionic acid was isolated in a yield of 1.40 g.

Compound 8: Melting point: 165-167°C. Combustion analysis: %C: 63.79 (calculated), 63.58 (found); %H: 4.98 (calculated), 4.92 (found); %N: 4.65 (calculated), 4.57 (found). ¹H NMR Analysis: (d₆-DMSO): δ 12.4, (COOH); δ 12.00, (OH); δ 10.3, s, 1H, (NH); δ 7.95, (CH aromatic ring alpha to amide); δ 7.55, (CH aromatic

ring alpha to ether); δ 7.40, (CH aromatic ring para to amide); δ 6.91, (CH aromatic ring meta to ether, and CH's ortho and para to hydroxy); δ 4.13, (CH₂ alpha to ether); δ 2.65, (CH₂ alpha to carboxylic acid).

5

Preparation of Compound 9

Compound 9 was prepared by the procedure described for the preparation of compound 8, however step B was omitted and O-anisoyl chloride was used in place of acetylsalicyloyl chloride.

10

Compound 9: Melting point: 154-158°C. Combustion analysis: %C: 64.76 (calculated), 64.70 (found); %H: 5.40 (calculated), 5.51 (found); %N: 4.44 (calculated), 4.24 (found). ¹H NMR Analysis: (d₆-DMSO): δ 12.4, (COOH); δ 10.0, s, 1H, (NH); δ 7.61, (CH aromatic ring alpha to amide, and CH's aromatic ring alpha to ether); δ 7.45, (CH aromatic ring meta to methoxy); δ 7.15, d, (CH aromatic ring alpha to methoxy); δ 7.03, (CH aromatic ring para to ether); δ 4.10, (CH₂ alpha to ether); δ 3.85, (CH₃); δ 2.65, (CH₂ alpha to carboxylic acid).

15
20

Preparation of Compound 7

Compound 7 was prepared by the procedure described for the preparation of compound 8. however step B was omitted and 2,3-dimethoxybenzoyl chloride was used in place of acetylsalicyloyl chloride.

25

Compound 7: Melting point: 135-137°C. Combustion analysis: %C: 62.61 (calculated), 62.31 (found); %H: 5.51 (calculated), 5.57 (found); %N: 4.06 (calculated), 3.95 (found). ¹H NMR Analysis: (d₆-DMSO): δ 12.3, (COOH); δ 10.1, (NH); δ 7.60, (CH's aromatic

30

ring alpha to ether); δ 7.09, (CH's aromatic ring ortho, meta and para to the 3-methoxy group); δ 6.88, dd, 2H, (CH aromatic ring meta to ether); δ 4.13, (CH₂ alpha to ether); δ 3.85, (CH₃ 2-methoxy group); δ 3.78, s, 3H, (CH₃ 3-methoxy group); δ 2.66, t, 2H, (CH₂ alpha to carboxylic acid).

Preparation of Compound 10

3-(4-Hydroxyphenyl)-1-propanol (20.0 g, 0.135 mole) and potassium carbonate (22.2 g, 0.162 mole) were stirred in acetone/water (150 mL/ 15 mL) in a 500mL round bottom flask equipped with a condenser, oil bath and a magnetic stirrer. Ethyl bromoacetate was added dropwise and the flask was heated to reflux for 18 hours. The reaction was cooled to room temperature and the reaction concentrated *in vacuo*. The residue was dissolved in ethyl acetate and washed with water, 1N hydrochloric acid and water again. The organics were dried over sodium sulfate, filtered and concentrated *in vacuo*. The structure of intermediate compound I, was confirmed by ¹H NMR and used without further purification, 30.29 g, 94.3% yield.

Intermediate compound, I (30.29 g, 0.127 mole) was dissolved in methylene chloride (100 mL) in a round bottom flask equipped with an addition funnel and an argon inlet/bubbler. The solution was cooled in an ice bath. Triethylamine (19.47 mL, 0.139 mole) was added dropwise over 5 minutes. Tosyl chloride (21.26 g, 0.127 mole) was dissolved in methylene chloride and added dropwise over 20 minutes to the reaction flask. The reaction was allowed to warm to room temperature and monitored by HPLC. Triethylamine and Tosyl chloride were added and the reaction monitored to completion. The mixture was concentrated

in vacuo and the residue dissolved in ethyl acetate. The solution was washed with hydrochloric acid (2 x 100 mL, 1 N) and saturated sodium bicarbonate (6 x 100 mL). The organics were dried over sodium sulfate and concentrated in vacuo. The structure of intermediate compound II, was confirmed by ¹H NMR and used without further purification, 29.15g, 60% yield.

In a 1 neck, 250 mL flask carsalam (4.22 g, 0.026 mole), intermediate compound II (10.0 g, 0.026 mole), N, N-dimethylacetamide (100 mL), and sodium carbonate (3.03 g, 0.0286 mole) were heated to 90°C under nitrogen for 20 hours. The reaction was cooled to room temperature and filtered to remove inorganics. Water (100 mL) was added and the solution extracted with ethyl acetate (3 x 100 mL). The organics were combined and concentrated in vacuo. The residue was stirred in sodium hydroxide (2N) and 1,4-dioxane for 2.5 hours and then acidified with concentrated hydrochloric acid. The resulting solids were recovered by filtration and dried in vacuo over phosphorous pentoxide to yield compound 10..

Compound 10: Melting point: 124-126.5°C. Combustion analysis for C₁₈H₁₉O₅N: %C: 65.64 (calculated), 65.32 (found); % H 5.81 (calculated), 5.89 (found); % N 4.25 (calculated), 4.15 (found).

¹H NMR Analysis: (300MHz, d₆-DMSO): δ..13.0; 8.85; 7.85; 7.40; 7.15; 6.85; 4.60; 3.30; 2.6; 1.8.

Preparation of the Ethanol Solvate of the Disodium Salt of Compound 1

In a 2L 4-neck flask equipped with condenser, thermocouple and recorder, variac (variable transformer), heating mantle, motorized stirrer, and addition funnel, 2-amino-4-chlorophenol

(200 g, 1.30 mole, available from Aldrich Chemical Company, Inc. of Milwaukee, Wisconsin, was stirred in water (500 ml). Acetic anhydride (171 g, 1.68 mole) was added to the reaction flask over 10 minutes. The reaction was heated to reflux for 30 minutes and then cooled to room temperature. The resulting slurry was vacuum filtered and solids washed with water yielding brown solids which were dried *in vacuo* at 50°C (249.2 g of N-acetyl-2-amino-4-chlorophenol; 97% yield; 95.4% by gas chromatography (GC)).

In a 2L, 4-neck flask equipped with condenser, thermocouple and recorder, nitrogen inlet/bubble, motorized stirrer, and addition funnel, ethanol (200mL) was charged. Sodium (9.5g, 0.412 mole) was added, after washing with kerosene, over 30 minutes keeping the reaction temperature below 60°C. The reaction was stirred until all of the sodium dissolved (about 90 minutes), while allowing the reaction to cool to 25°C. 61.25 g (0.33 mole) N-acetyl-2-amino-4-chlorophenol was dissolved in ethanol (350 mL) at 70°C and added to the sodium solution by cannula. Ethyl 8-bromooctanoate (103 g, 0.412 mole) (available from Riedel-de Haen, Seelze, Germany) was then added to the reaction by addition funnel. The reaction was heated to reflux for about 18.5 hours. After cooling to room temperature, the reaction was filtered and the filtrate condensed *in vacuo*. The resulting brown semi-solid was stirred with heptane (600 mL) for several hours, filtered and dried *in vacuo* yielding 120.8 g of 2-N-acetylamino-4-(chlorophenoxy)ethyl octanoate as a solid, 100% yield.

In a 5 liter, 4-neck flask equipped with a motorized stirrer, thermometer, THERM-O-WATCH® heat controller (available from I²R, Cheltenham, PA), heating mantle, and a nitrogen

inlet/bubbler, sodium hydroxide (240.0 g, 6.0 mole) was dissolved in ethanol (120 mL) at 46.5°C. 12.0 g (0.337 mole) 2-N-acetylamino-4-(chlorophenoxy)ethyl octanoate was added to the solution and the flask was heated to 80°C for about 13 hours.

5 The hydrolysis was followed by HPLC. Ethanol (347.8 mL) was distilled off and the reaction was cooled to 25°C producing solids. Concentrated hydrochloric acid (469.1 mL) was added to the flask dropwise to a pH of 4.65 keeping the temperature below 25°C with an ice bath. Brown solids were filtered off and washed
10 with water yielding 82.93 g (86% yield) 8-(2-amino-4chloro-phenoxy)octanoic acid. The structure was verified by NMR. A second batch of 81.33g of the same compound was made by the same method.

In a 3 liter 4-neck flask equipped with motorized stirrer,
15 ice bath, addition funnel, condenser and thermometer, dry methanol (500 mL) was cooled to 2°C. Acetyl chloride (52.56 g, 0.669 mole) was added dropwise to the methanol and the solution was stirred for an additional 15 minutes. 159 g (0.558 mole) 8-(2-amino-4-chloro-phenoxy)octanoic acid was dissolved in dry
20 methanol (500 mL) and slowly added to the reaction flask. After stirring for 15 minutes the ice bath was removed and the flask heated to reflux for 3 hours. The solution was concentrated in vacuo yielding 187 g (100%) of methyl (8-(2-amino-4-chloro-phenoxy)octanoate•HCl) as a light brown solid. The structure was
25 confirmed by NMR.

In a 3 liter 4-neck flask equipped with motorized stirrer, addition funnel and thermometer, 100 g (0.332 mole) methyl (8-(2-amino-4-chloro-phenoxy)octanoate•HCl) was stirred in THF (500 mL) and triethylamine (70.55 g, 0.697 mole) was added dropwise
30 at 5°C. Acetylsalicyloyl chloride (65.98 g, 0.332 mole)

dissolved in THF (250mL) was then added to the reaction over 30 minutes. After an additional 15 minutes at 5°C the reaction was warmed to room temperature and monitored by HPLC. Sodium hydroxide (2N, a total of 950 mL) was added to the reaction flask and the hydrolysis followed by HPLC. The reaction was cooled to 5°C with an ice bath and concentrated hydrochloric acid (170mL) was added dropwise to the reaction keeping the temperature below 20°C to a final pH of 4.0. The THF was distilled off and the resulting tan solid filtered off and air-dried, yielding 117.91 g (87.5% yield) of 8-[(N-Salicyloyl-2-amino-4-chloro)phenoxy]octanoic acid (Compound III). The structure was confirmed by NMR and elemental analysis (calculated for $C_{21}H_{24}NO_5Cl$: C, 62.14; H, 5.96; N, 3.45; analyzed: C, 61.92; H, 5.91; N, 3.26).

In a 2 liter 3-neck flask equipped with a motorized stirrer, heating mantel, thermometer, nitrogen inlet/bubbler and addition funnel, 117.91 g Compound III was stirred in ethanol (1L) and heated to 75°C for 30 minutes. The temperature was lowered to 70°C and sodium hydroxide in ethanol (11.3wt%, 199.7 mL) was added to the solution and stirred for 30 minutes. Ethanol (970 mL) was distilled off and the reaction cooled slightly. The molten mass was poured into a crystallizing dish and dried at 80°C, full vacuum grinding of solids after they formed, yielding (130.36 g (97.4%) disodium 8-(N-salicyloyl-2-amino-4-chloro)phenoxy)octanoate, ethanol solvate (Compound I ethanol solvate). The structure of the ethanol solvate was verified by elemental analysis: calculated for $C_{23}H_{28}NO_6Na_2Cl \cdot 1.3664 H_2O$: C, 55.31; H, 5.19; N, 2.80; Na, 9.21; found: C, 54.4; H, 5.42; N, 2.90; Na, 7.98. The structure of the ethanol solvate was verified by 1H NMR (300 MHz, DMSO- d_6): δ

14.8, 8.70, 7.7, 6.9, 6.55, 6.2, 4.0, 4.4 3.45, 1.4, 1.24.4 ,
ethanol; 4.4 , 3.45 , 1.05. Melting point >250°C (limit of
instrument used).

5 **Example 2**

Pulmonary Insulin

Dosing compositions of delivery agent compound and human
insulin in water were prepared. Typically, deionized water was
added to 1.5 mg of delivery agent compound to bring the volume
10 to 1.0 ml, and the solution was vortexed. Either the sodium
salt of the delivery agent compound was used or the free acid
was converted to the sodium salt by stirring the resultant
solution and adding one equivalent of sodium hydroxide (10 N)
and diluting with water. The solution was vortexed, then heated
15 (about 37°C) and sonicated. The pH was adjusted to between about
7.0 to 8.5 with NaOH or HCl.

75 µl human insulin stock solution (2 mg/ml) was added to
the solution. The human insulin stock solution was prepared as
follows. To 0.02 g human insulin was added 3 ml HCl solution
20 (pH = 3) in deionized water, the pH of the resulting solution
was brought to about 2.6 with HCl and NaOH was added until the
solution was clear. The pH was then raised to 7.6 using NaOH
and HCl. The final volume was brought to 10 ml with pH 7.5
deionized water so that the final pH was 7.59. Water was added
25 to bring the total volume to 2.0 ml, and the solution was gently
inverted several times. The final delivery agent compound dose,
insulin dose and volume dose amounts are as listed below in the
tables.

The typical dosing and sampling protocols were as follows.
30 Male Sprague-Dawley rats weighing between 200-250g were fasted
for 24 hours and administered ketamine (44 mg/kg) and

chlorpromazine (1.5 mg/kg) 15 minutes prior to dosing.

Typically, a dosing group of five rats was administered one of the dosing solutions. A control group of five animals was dosed with insulin alone. A tracheal instillator for rodents,
5 equipped with light (available from Penn Century, Inc., Pittsburgh, PA) was filled with dosing solution and inserted down the throat until the needle went into the trachea (confirmed visually). Solution was administered by pressing the plunger.

10 Blood samples were collected serially from the tail artery, typically at time = 5, 15, 30, 60 and 120 minutes. Serum insulin levels were determined with an Insulin ELISA Test Kit (Kit # DSL-10-1600 from Diagnostic Systems Laboratories, Inc., Webster, TX). The results from the five rats in each dosing
15 group were averaged for each time point. The maximum is reported below in Table 1.

Insulin - Oral Delivery

Oral dosing (PO) compositions of delivery agent compound
20 and human zinc insulin (minimum 26 IU/mg available from Calbiochem - Novabiochem Corp, La Jolla, CA) were prepared in deionized water. Typically, 500 mg of delivery agent compound was added to 1.5 ml of water. The free acid of the delivery agent compound was converted to the sodium salt by stirring the
25 resultant solution and adding one equivalent of sodium hydroxide. The solution was vortexed, then heated (about 37°C) and sonicated. The pH was adjusted to about 7 to 8.5 with NaOH or HCl. Additional NaOH was added, if necessary, to achieve uniform solubility, and the pH re-adjusted to about 7 to 8.5.
30 Water was then added to bring the total volume to about 2.4 ml and vortexed. About 1.25 mg insulin from an insulin stock

solution (15 mg/ml made from 0.5409 g insulin and 18 ml deionized water, adjusting with HCl and NaOH to obtain a clear solution using 40 ml concentrated HCl, 25 ml 10N NaOH and 50 ml 1N NaOH) was added to the solution and mixed by inverting. The solution was used in the dosing protocol immediately, or alternatively, the solution was placed in a 37°C water bath for one hour prior to dosing. The final delivery agent compound dose, insulin dose and dose volume amounts are listed below in Table 1.

The typical dosing and sampling protocols were as follows. Male Sprague-Dawley rats weighing between about 200-250g were fasted for 24 hours and administered ketamine (44 mg/kg) and chlorpromazine (1.5 mg/kg) 15 minutes prior to dosing and again as needed to maintain anesthesia. A dosing group of five rats was administered one of the dosing solutions. For oral dosing, an 11 cm Rusch 8 French catheter was adapted to a 1 ml syringe with a pipette tip. The syringe was filled with dosing solution by drawing the solution through the catheter, which was then wiped dry. The catheter was placed down the esophagus leaving 1 cm of tubing past the incisors. The dosing solution was administered by pressing the syringe plunger.

Blood samples were collected serially from the tail artery, typically at time = 15, 30, 60, 120 and 180 minutes. Serum insulin levels were determined with an Insulin ELISA Test Kit (Kit # DSL-10-1600 from Diagnostic Systems Laboratories, Inc., Webster, TX), modifying the standard protocol in order to optimize the sensitivity and linear range of the standard curve for the volumes and concentrations of the samples used in the test kit. Serum human insulin concentrations (μ U/ml) were measured for each time point for each of the five rats in each dosing group. The five values for each time point were

averaged and the results plotted as serum insulin concentration versus time. The maximum of the averaged values is reported below in Table 1. Previous experiments revealed no measurable levels of human insulin following oral dosing with human insulin alone.

Table 1. Pulmonary/Oral Delivery of Insulin

Compound	Method of Administration	Volume Dose (ml/kg)	Compound Dose (mg/kg)	Insulin Dose (mg/kg)	Mean Peak Serum Insulin (insulin + Compound) (pg/ml \pm SD) (SE)
1	Pulmonary	0.4	0.3	0.03	73.19 \pm 41.6
1	Pulmonary	0.4	0.06	0.03	156.04 \pm 87.02
1	Pulmonary	0.4	0.15	0.03	119.89 \pm 29.57
1	Pulmonary	0.4	0.3	0.03	151.50 \pm 84.63
1	Pulmonary	0.4	0.06	0.03	200.27 \pm 87.14
1	Pulmonary	0.4	0.15	0.03	185.09 \pm 128.11
1	Pulmonary	0.4	0.3	0.03	129.44 \pm 17.49
1	Pulmonary	0.4	0.06	0.03	73.37 \pm 19.93
1	Pulmonary	0.4	0.03	0.03	181.80 \pm 197.67
1	Pulmonary	0.4	0.15	0.03	93.56 \pm 49.84
1	Pulmonary	0.4	0.06	0.03	122.35 \pm 96.64
1	Pulmonary	0.4	0.03	0.03	103.73 \pm 33.96
1	Pulmonary	0.4	0.07	0.03	128.10 \pm 36.31
1	Pulmonary	0.4	0.3	0.03	123.08 \pm 40.28
1	Pulmonary	0.4	0.3	0.03	87.02 \pm 26.09
2	Pulmonary	0.4	0.3	0.03	11.6 \pm 2.27
2	PO	1	200	0.5	5.9 \pm 7.94
3	PO	1	200	0.5	110.59 \pm 11.84
4	PO	1	200	0.5	6.56 \pm 9.22
4	PO	1	200	0.5	5.90 \pm 7.90
5	PO	1	200	0.5	45.45 \pm 74.04
5	PO	1	200	0.5	69.44 \pm 62.14

10 Salmon Calcitonin (sCT)

Oral Delivery

Oral dosing (PO) compositions of delivery agent compound and salmon calcitonin (sCT) in water were prepared. Typically,

450 mg of delivery agent compound was added to 2.0 ml of water.

Either the sodium salt of the delivery agent compound was used or the free acid was converted to the sodium salt by stirring the resultant solution and adding one equivalent of sodium

5 hydroxide (1.0 N) and diluting with water. The solution was vortexed, then heated (about 37°C) and sonicated. The pH was adjusted to about 7 (6.5 to 8.5) with NaOH or HCl. 90 µg sCT from a stock solution was added to the solution. Water was then added to bring the total volume to about 3.0 ml (varied
10 depending on the solubility of the delivery agent compound) and vortexed. The final delivery agent compound dose, sCT dose and volume dose amounts are listed below in Table 2.

The typical dosing and sampling protocols were as follows. Male Sprague-Dawley rats weighing between 200-250g were fasted
15 for 24 hours and administered ketamine (44 mg/kg) and chlorpromazine (1.5 mg/kg) 15 minutes prior to dosing. A dosing group of five rats was administered one of the dosing solutions.

For oral dosing, an 11 cm Rusch 8 French catheter was adapted to a 1 ml syringe with a pipette tip. The syringe was filled
20 with dosing solution by drawing the solution through the catheter, which was then wiped dry. The catheter was placed down the esophagus leaving 1 cm of tubing past the rat's incisors. The dosing solution was administered by pressing the syringe plunger.

25 Blood samples were collected serially from the tail artery, typically at time = 0, 10, 20, 30, 60 and 90 minutes. Serum sCT levels were determined by testing with an EIA kit (Kit # EIAS-6003 from Peninsula Laboratories, Inc., San Carlos, CA). The observed sCT levels were adjusted according to baseline values
30 obtained at time = 0. The results from the five rats in each dosing group were averaged for each time point. The maximum is

reported below in Table 2.

Table 2. Oral Delivery of Salmon Calcitonin (sCT)

Compound	Volume Dose (ml/kg)	Compound Dose (mg/kg)	sCT Dose (μ g/kg)	Mean Peak Serum Sct (pg/ml \pm SD)
1	1	150	30	164 \pm 156
2	1	150	30	112 \pm 155
2	1	150	30	482 \pm 389
2	1	150	30	206 \pm 461
7	1	150	30	206 \pm 138

5 **Preparation of the Disodium Salt of Compound 1**

25g of the ethanol solvate of compound prepared in Example 1 was dissolved in 250 ml methanol. 5 g of activated carbon was added and the mixture was stirred for 20 minutes at 50-60° C. The slurry was filtered and the carbon treatment was repeated.

10 Methanol was removed *in vacuo*. The resulting solid was washed with 100 ml of hexane and dried *in vacuo*. 14.8 g of the disodium salt of compound 1 was recovered as an off-white solid. ¹H NMR indicated that there was no ethanol associated with the purified sample.

15 The disodium salt of compound 1 was tested *in vivo* with insulin by the procedure described above for oral administration of insulin. The results are shown below in Table 3.

Table 3

Pulmonary Delivery of Insulin with the Disodium Salt of Compound I

Compound	Volume Dose (ml/kg)	Compound Dose (mg/kg)	Insulin Dose (mg/kg)	Mean Peak Serum Insulin (insulin + Compound) (pg/ml \pm SD) (SE)
1	0.4	0.36	0.03	166.97 \pm 9.64
1	0.4	0.18	0.03	172.10 \pm 36.31

Compound	Volume Dose (ml/kg)	Compound Dose (mg/kg)	Insulin Dose (mg/kg)	Mean Peak Serum Insulin (insulin + Compound) (pg/ml \pm SD) (SE)
1	0.4	0.06	0.03	72.36 \pm 29.76
1	0.4	0.15	0.03	35.96 \pm 8.5
1	0.4	0.03	0.03	50.66 \pm 19.13

Heparin Delivery Intracolonic Delivery

Intracolonic (IC) dosing solutions containing a delivery agent compound and heparin sodium USP in 25% aqueous propylene glycol were prepared. Either the sodium salt of the delivery agent compound was used or the free acid was converted to the sodium salt with one equivalent of sodium hydroxide. Typically, the delivery agent compound and heparin (about 166-182 IU/mg) were mixed by vortex as dry powders. This dry mixture was dissolved in 25% v/v aqueous propylene glycol, vortexed and placed in a sonicator (about 37°C). The pH was adjusted to about 7 (6.5 to 8.5) with aqueous NaOH (2N). The dosing solution was sonicated to produce a clear solution. The final volume was adjusted to 3.0 mL. The final delivery agent compound dose, heparin dose and volume dose amounts are shown in Table 4, below.

The typical dosing and sampling protocols were as follows. Male Sprague-Dawley rats weighing between 275-350g were fasted for 24 hours and were anesthetized with ketamine hydrochloride (88 mg/kg) intramuscularly immediately prior to dosing. A dosing group of five rats was administered one of the dosing solutions.

For intracolonic (IC) dosing, a 7.5 cm 8 fr Rusch catheter was adapted to a 1 ml syringe with a pipette tip. The dosing catheter was inserted into the colon through the anus until the tube was no longer visible. The dosing solution was expressed

slowly into the colon.

Citrated blood samples were collected by cardiac puncture following the administration of ketamine (88 mg/kg), typically at time - 0.25, 0.5, 1.0 and 1.5 hours. Heparin activity was
 5 determined by utilizing the activated partial thromboplastin time (APTT) according to the method of Henry, J.B., Clinical Diagnosis and Management by Laboratory Methods, Philadelphia, PA, W.B. Saunders (1979). Previous studies indicated baseline values of about 20 seconds. Results from the five rats in each
 10 group were averaged for each time point. The maximum is reported below in Table 4.

Table 4. Intracolonic Delivery of Sodium Heparin USP in Rats

Compound	Volume Dose (ml/kg)	Compound Dose (mg/kg)	Insulin Dose (mg/kg)	Mean Peak Serum Insulin (insulin + Compound) (pg/ml \pm SD)
1	1	750	50	0
2	1	50	25	74.12 \pm 126.28
2	1	50	25	269 \pm 42
3	1	50	25	18.3 \pm 2.6
8	1	20	100	262.2 \pm 182.8
9	1	50	25	68.2 \pm 37.1

Parathyroid Hormone Delivery (PTH 1-34) Oral/ Intracolonic Delivery

Oral gavage (PO) dosing solutions of delivery agent
 25 compound and human parathyroid hormone residues 1-34 (PTH) in water were prepared. A solution of the delivery agent compound was made either with the sodium salt of the delivery agent compound or by converting the free acid to its sodium salt. Typically, a solution of the delivery agent compound was
 30 prepared in water and stirred, adding one equivalent of sodium hydroxide (1.0 N) when making the sodium salt. The final dosing

solutions were prepared by mixing the solution of the delivery agent compound with a PTH stock solution (typically having a concentration of 5 mg PTH/ml) and diluting to the desired volume (usually 3.0 ml). The delivery agent compound, PTH and volume dose amounts are shown in Table 5 below.

The typical dosing and sampling protocols were as follows. Male Sprague-Dawley rats weighing between 200-250g were fasted for 24 hours and administered ketamine (44 mg/kg) and chlorpromazine (1.5 mg/kg) 15 minutes prior to dosing. A dosing group of five rats was administered one of the dosing solutions.

For oral gavage (PO) dosing, an 11 cm Rusch 8 French catheter was adapted to a 1 mL syringe with a pipette tip. The syringe was filled with dosing solution by drawing the solution through the catheter, which was then wiped dry. The catheter was placed down the esophagus leaving 1 cm of tubing past the rat's incisors. The dosing solution was administered by pressing the syringe plunger.

Blood samples were collected serially from the tail artery, typically at time = 0, 15, 30, 45, 60 and 90 minutes for oral dosing. Serum PTH concentrations were quantified by a PTH radioimmunoassay kit (Kit # RIK 6101 from Peninsula Laboratories, Inc. San Carlos, CA). Previous studies indicated baseline values of about zero. Results from the five rats in each group were averaged for each time point. The maximum is reported below in Table 5.

Table 5. Oral Delivery of PTH in Rats

Compound	volume dose (ml/kg)	Compound Dose (mg/kg)	PTH Dose (mg/kg)	Mean Peak Serum [PTH] (pg/ml) + SD
8	1	100	200	767.21 + 140.35
8	1	100	200	276.31 + 330.06

8	1	100	200	11.31 + 25.24
---	---	-----	-----	---------------

Interferon - Oral Delivery

Dosing solutions of delivery agent compound and human interferon (IFN) were prepared in deionized water. The free acid of the delivery agent compound was converted to the sodium salt with one equivalent of sodium hydroxide. Typically, a solution of the delivery agent compound was prepared in water and stirred, adding one equivalent of sodium hydroxide (1.0 N) when making the sodium salt. This mixture was vortexed and placed in a sonicator (about 37°C). The pH was adjusted to about 7.0 to 8.5 with aqueous NaOH. The mixture was vortexed to produce a uniform suspension or solution, also using sonication and heat if necessary. Additional NaOH was added, if necessary, to achieve uniform solubility, and the pH re-adjusted to about 7.0 to 8.5. The delivery agent compound solution was mixed with an IFN stock solution (about 22.0 to 27.5 mg/ml in phosphate buffered saline) and diluted to the desired volume (usually 3.0 ml). The final delivery agent compound and IFN doses, and the dose volumes are shown below in Table 6.

The typical dosing and sampling protocols were as follows. Male Sprague-Dawley rats weighing between 200-250g were fasted for 24 hours and administered ketamine (44 mg/kg) and chlorpromazine (1.5 mg/kg) 15 minutes prior to dosing and again as needed to maintain anesthesia. A dosing group of five rats was administered one of the dosing solutions. An 11cm Rusch 8 French catheter was adapted to a 1 ml syringe with a pipette tip. The syringe was filled with dosing solution by drawing the solution through the catheter, which was then wiped dry. The catheter was placed down the esophagus leaving 1 cm of tubing past the incisors. The dosing solution was administered

by pressing the syringe plunger.

Blood samples were collected serially from the tail artery, typically at time = 0, 15, 30, 45, 60 and 90 minutes. Serum IFN concentrations were determined with a Cytoscreen Immunoassay Kit
5 for human IFN-alpha (catalog # KHC4012 from Biosource International, Camarillo, CA). Previous studies indicated baseline values of about zero. Results from the rats in each group were averaged for each time point. The maximum of these averages (i.e., the mean peak serum IFN concentration) is
10 reported below in Table 6.

Table 6. Interferon - Oral Delivery

Delivery Agent Compound	Volume dose (ml/kg)	Delivery Agent Compound Dose (mg/kg)	IFN Dose (mg/kg)	Mean Peak Serum [IFN] (ng/ml) \pm SD
10	1	200	2	0.1256 \pm 0.280

15 **Recombinant Human Growth Hormone (rhGH) Oral Delivery**

Oral gavage (PO) dosing solutions of delivery agent compound and rhGH in phosphate buffer were prepared. A solution of the delivery agent compound was made either with the sodium salt of the compound or by converting the free acid to its
20 sodium salt. Typically, a solution of the delivery agent compound was prepared in phosphate buffer and stirred, adding one equivalent of sodium hydroxide (1.0 N) when making the sodium salt. The final dosing solutions were prepared by mixing the solution containing the delivery agent compound with an rhGH
25 stock solution (15 mg rhGH/ml) and diluting to the desired volume (usually 3.0 ml). The delivery agent compounds and rhGH dose amounts are shown in Table 7 below.

The typical dosing and sampling protocols were as follows. Male Sprague-Dawley rats weighing between 200-250g were fasted for 24 hours and administered ketamine (44 mg/kg) and chlorpromazine (1.5 mg/kg) 15 minutes prior to dosing. A dosing group of five rats was administered one of the dosing solutions. For oral gavage (PO) dosing, an 11 cm Rusch 8 French catheter was adapted to a 1 mL syringe with a pipette tip. The syringe was filled with dosing solution by drawing the solution through the catheter, which was then wiped dry. The catheter was placed down the esophagus leaving 1 cm of tubing past the rat's incisors. The dosing solution was administered by pressing the syringe plunger

Blood samples were collected serially from the tail artery, typically at time = 0, 15, 30, 45, 60 and 90 minutes for oral and 0, 10, 20, 30, 60 and 90 for IC dosing. The five samples from each time period were pooled. Serum rhGH concentrations were determined with an rhGH immunoassay test kit (Kit #K1F4015 from Genzyme Corporation Inc., Cambridge, MA). Previous studies indicated baseline values of about zero.

The maximum concentration for each group is reported below in Table 7.

Table 7. Oral Delivery of rhGH in Rats

Compound	Volume dose (ml/kg)	Compound Dose (mg/kg)	rhGH Dose (mg/kg)	Mean Peak Serum (ng/ml)
7	1	300	3	49.55
9	1	200	3	0

The above-mentioned patents, applications, test methods, and publications are hereby incorporated by reference in their entirety.


Many variations of the present invention will suggest themselves to those skilled in the art in light of the above detailed description. All such obvious variations are within the fully intended scope of the appended claims.

5

WHAT IS CLAIMED IS:

1. A compound of the formula

5



Formula I

wherein:

- 10 R^1 , R^2 , R^3 , R^4 , and R^5 are independently selected from hydrogen, halogen, $-\text{OH}$, $-\text{OCH}_3$, C_1 - C_4 alkyl, $-\text{NH}_2$, $-\text{NH}(\text{CH}_3)$, $-\text{N}(\text{CH}_3)_2$, or $-\text{NO}_2$;

m is an integer ranging from 0 to 4;


- 15 R^6 is a phenyl substituted with $-\text{O}-\text{R}^7-\text{COOH}$ at the ortho, meta, or para position;

R^6 is optionally substituted with one or more members selected from halogen, $-\text{OH}$, $-\text{OCH}_3$, C_1 - C_4 alkyl, $-\text{NH}_2$, $\text{NH}(\text{CH}_3)$, $-\text{N}(\text{CH}_3)_2$, and $-\text{NO}_2$; and


- 20 R^7 is C_1 - C_{12} alkyl, or a salt thereof.

2. The compound of claim 1 wherein R^1 is $-\text{OH}$


3. The compound of claim 1 wherein said compound is
25 selected from the group consisting of:




Compound 1




Compound 2




Compound 3



Compound 4




Compound 5

Compound 6

and the salts and solvates of said compounds.

5

4. The compound of claim 1 wherein said compound is



Compound 1

10



and the salts and solvates of said compounds.

5. The compound of claim 4 wherein said compound is the disodium salt.


15

6. The compound of claim 1 wherein said compound is selected from the group consisting of:


20

Compound 7

5

Compound 8

10

Compound 9

15

Compound 10

and the salts and solvates of said compounds.

7. A composition comprising:
 (A) an active agent; and
 (B) a compound as defined in claim 1.

- 5 8. A composition comprising:
 (A) an active agent; and
 (B) a compound as defined in claim 4.

- 10 9. A composition comprising:
 (A) an active agent; and
 (B) a compound as defined in claim 5.

15 10. The composition of **claim 7**, wherein the active agent is selected from the group consisting of a biologically active agent, a chemically active agent, and a combination thereof.

20 11. The composition of **claim 10**, wherein the biologically active agent comprises at least one protein, polypeptide, peptide, hormone, polysaccharide, mucopolysaccharide, carbohydrate, or lipid.

25 12. The composition of **claim 11**, wherein the biologically active agent is selected from the group consisting of: growth hormones, including but not limited to human growth hormones, recombinant human growth hormones, bovine growth hormones, porcine growth hormones; growth hormone releasing hormones; growth hormone releasing factor; interferons, including but not limited to α -interferon, β -interferon, γ -interferon; interleukin including but not limited to
30 interleukin-1, and interleukin-2; insulin including but not limited to, porcine insulin, bovine insulin, human insulin,

human recombinant insulin, insulin-like growth factor, insulin-like growth factor -1; heparin including but not limited to unfractionated heparin, heparinoids, dermatans, chondroitins, low molecular weight heparin, very low molecular weight heparin, ultra low molecular weight heparin; calcitonin including but not limited to salmon calcitonin, eel calcitonin, human calcitonin; erythropoietin; atrial natriuretic factor; antigens; monoclonal antibodies; somatostatin; protease inhibitors; adrenocorticotropin; gonadotropin releasing hormone; oxytocin; leutinizing-hormone-releasing-hormone; follicle stimulating hormone; glucocerebrosidase; thrombopoietin; filgrastim; prostaglandins; cyclosporin; vasopressin; cromolyn sodium; sodium chromoglycate; disodium chromoglycate; vancomycin; desferrioxamine; parathyroid hormone; fragments of parathyroid hormone; antimicrobials; anti-fungal agents; vitamins; analogs, fragments, mimetics and polyethylene glycol (PEG)-modified derivatives of these compounds; and any combination thereof.

13. The composition of **claim 12**, wherein the biologically active agent comprises insulin, unfractionated heparin, low molecular weight heparin, very low molecular weight heparin, ultra low molecular weight heparin, calcitonin, PTH, EPO, hGH or combinations thereof.

14. The composition of **claim 13**, wherein the biologically active agent comprises insulin.

15. A dosage unit form comprising:

(A) the composition of **claim 7**; and

- (B) (a) an excipient
(b) a diluent,
(c) a disintegrant,
(d) a lubricant,
5 (e) a plasticizer,
(f) a colorant,
(g) a dosing vehicle, or
(h) any combination thereof.

10 16. The dosage unit form of **claim 15**, wherein the active agent is selected from the group consisting of a biologically active agent, a chemically active agent, and a combination thereof.

15 17. The dosage unit form of **claim 16**, wherein the biologically active agent comprises at least one protein, polypeptide, peptide, hormone, polysaccharide, mucopolysaccharide, carbohydrate, or lipid.

20 18. The dosage unit form of **claim 17**, wherein the biologically active agent is selected from the group consisting of: : growth hormones, including but not limited to human growth hormones, recombinant human growth hormones, bovine growth hormones, porcine growth hormones; growth hormone releasing
25 hormones; growth hormone releasing factor; interferons, including but not limited to α -interferon, β -interferon, γ -interferon; interleukin including but not limited to interleukin-1, and interleukin-2; insulin including but not limited to, porcine insulin, bovine insulin, human insulin,
30 human recombinant insulin, insulin-like growth factor, insulin-like growth factor -1; heparin including but not limited to

unfractionated heparin, heparinoids, dermatans, chondroitins,
low molecular weight heparin, very low molecular weight heparin,
ultra low molecular weight heparin; calcitonin including but not
limited to salmon calcitonin, eel calcitonin, human calcitonin;
5 erythropoietin; atrial naturetic factor; antigens; monoclonal
antibodies; somatostatin; protease inhibitors;
adrenocorticotropin; gonadotropin releasing hormone; oxytocin;
leutinizing-hormone-releasing-hormone,;follicle stimulating
hormone; glucocerebrosidase; thrombopoietin; filgrastim;
10 prostaglandins; cyclosporin; vasopressin; cromolyn sodium;
sodium chromoglycat; disodium chromoglycate; vancomycin;
desferrioxamine; parathyroid hormone; fragments of parathyroid
hormone; antimicrobials; anti-fungal agents; vitamins; analogs,
fragments, mimetics and polyethylene glycol (PEG)-modified
15 derivatives of these compounds; and any combination thereof.

19. The dosage unit form of **claim 18**, wherein the
biologically active agent comprises insulin, unfractionated
heparin, low molecular weight heparin, very low molecular weight
20 heparin, ultra low molecular weight heparin, calcitonin, PTH,
EPO, hGH, or combinations thereof.

20. The dosage unit form of **claim 19**, wherein the active
agent comprises insulin.

25

21. The dosage unit form of **claim 15**, wherein the dosage
unit form is a tablet, a capsule, a powder, or a liquid.

22. The dosage unit form of **claim 15**, wherein dosing unit
30 form comprises a dosing vehicle and dosing vehicle is selected
from the group consisting of water, 25% aqueous propylene

glycol, phosphate buffer, 1,2-propane diol, ethanol, and any combination thereof.

23. A method for administering a biologically-active agent
5 to an animal in need of the agent, the method comprising
administering to the animal the composition of **claim 7**.

24. A method for administering a biologically-active agent
to an animal in need of the agent, the method comprising
10 administering to the animal via the pulmonary route the
composition of **claim 9**.

25. A method for preparing a composition comprising
mixing:
15 (A) at least one active agent;
(B) the compound of claim 1; and
(C) optionally, a dosing vehicle.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US01/01274

A. CLASSIFICATION OF SUBJECT MATTER IPC(7) : C07C 229/00; A61K 38/28; A61K 51/00 US CL : 562/455; 514/3, 424/1.11 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 562/455; 514/3, 424/1.11 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X — Y	BROWN et al. Receptor Binding Sites of Hypoglycemic Sulfonylureas and Related [(Acylamino)alkyl]benzoic Acids. J. Med. Chem. 01 March 1983, Vol. 27, pages 79-81, especially pages 79 and 80.	1, 7, 23 ----- 2, 3, 4, 5, 6, 8, 9, 10, 11, 12, 13-22, 25
X — Y	GOA et al. Bezafibrate. An Update of its Use in the Management of Dyslipidaemia. Drugs, November 1996, Vol. 52 (5), pages 726-753, especially pages 733 and 745.	1, 7, 23, 25 ----- 2-6; 8-22
X	WO 98/05331(PATERNITI ET AL) 12 February 1998 (2/02/98), see the entire document	1-23, 25
X	WO 99/40904(BONHOMME ET AL) 19 August 1999 (19/08/99), see the entire document.	1- 23, 25
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family	
Date of the actual completion of the international search 13 APRIL 2001		Date of mailing of the international search report 04 MAY 2001
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer HECTOR M. REYES Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US01/01274

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 5,122, 539(ABRAHAM ET AL) 16 June 1992 (16/06/92), see the entire document, especially fig. 8, compounds 1-7 and 32.	1
X	US 5,304,575 (W. BECK) 19 April 1994 (19/04/94), see the entire document, especially col. 2, lines 3 to 11.	1, 7, 10, 11, 12, 14, 16, 21, 23, 25